## NIEMANN-PICK TYPE C DISEASE (NP-C) IS A

## ARE PROGRESSIVE IRREVERSIBLE C CHRONICALLY DEBILITATING YSOSOMAL STORAGE DISEASE







Incidence of NP-C is 1 in 90.000 live births4

Likely an underestimate due to lack of clinical awareness1

**THINK AGAIN** THINK NP-C

THINK AGAIN. THINK NP-C aims to support healthcare professionals unfamiliar with NP-C to recognise the key signs and symptoms of NP-C and reduce the time to diagnosis

## Individual symptoms are non-specific to the disease<sup>1,3</sup>

If vou are a: Paediatrician

**LOOK FOR ATAXIA, DEVELOPMENTAL DELAY, HEPATOSPLENOMEGALY** 



Paediatric hepatologist/neonatologist

LOOK FOR HEPATO/SPLENOMEGALY, **NEONATAL CHOLESTATIC JAUNDICE,** NEONATAL LIVER DYSFUNCTION

Adult neurologist/psychiatrist

LOOK FOR COGNITIVE DECLINE. **ORGANIC PSYCHOSIS, PROGRESSIVE ATAXIA** 



- Patterson M, Hendriksz, Walterfang M, et al. on behalf of the NP-C Guidelines Working Group. Recommendations for the diagnosis and management of Niemann-Pick disease type C: an update. Mol Genet Metab 2012; 106(3): 330–344.
  Vanier, M. Niemann-Pick disease type C. Orphanet J Rare Dis 2010; 5: 16.
  Wijburg FA, Sedel F, Pineda M, et al. Development of a suspicion index to aid diagnosis of Niemann-Pick disease type C. Neurology 2012;78(20):1560–1567.

- Wassif C, Cross J, Iben J et al. High incidence of unrecognized visceral/neurological late-onset Niemann-Pick disease, type C1, predicted by analysis of massively parallel sequencing data sets. Genet Med 2016; 18(1): 41–48
- 5. Mengel E, Klünemann H, Lourenço C, et al. Niemann-Pick disease type C symptomatology: an expert-based clinical description. Orphanet J Rare Dis 2013; 8: 166.

**NP-C** takes on average 5 YEARS to diagnose





...waiting for an answer, watching a loved one get worse

Have you checked for eye movement abnormalities?





**Vertical supranuclear gaze** palsy (VSGP) is present in virtually all patients<sup>1,3</sup>

To help reduce the time to diagnosis visit www.think-npc.com today



